For further information about this service contact:





Children's Services Union Street

Bedford MK40 2SF

Tel: 01234 315847

Email: ccs.beds.childrens.cct@nhs.net

Information and guidance can be found at:

- ERIC Helpline
- Bladder & Bowel UK
- NHS Choices
- NICE

0845 370 8008 www.eric.org.uk 0161 607 8200 www.disabledliving.co.uk www.nhs.uk www.nice.org.uk/guidance

Continent

Confident

Carefree













If you require this information in a different format such as in large print or on audio tape, or in a different language please contact the service on the details above.

If you have any compliments about this service or suggestions for improvements, contact our Patient Advice and Liaison Service on 0300 131 1000 (charges may apply depending on your network) or email: ccs-tr.pals@nhs.net.

For free, confidential health advice and information 24 hours a day, 365 days a year please contact NHS 111.

©Cambridgeshire Community Services NHS Trust Website: www.cambscommunityservices.nhs.uk Code No: Date of Production: Date of Review:

0559 - V1.2 (Jan 2020) n: June 2018 June 2021

Enuresis Alarm - Simple Measures

Children's Continence Clinic



Bedfordshire adult and children's community health services are provided in partnership by East London NHS Foundation Trust (ELFT) and Cambridgeshire Community Services (CCS) NHS Trust Please ensure that you continue following the simple measures outlined below:

Drinks

- Your child should be drinking at least 6 8 cups of **water** (or squash) each day and even more in the warm weather or when taking part in sport.
- Reduce or stop all fizzy drinks, blackcurrant, orange juice, caffeine and hot chocolate.
- Stop drinks two hours before bedtime.
- Ensure drinks are taken regularly throughout the day.

Toileting

- Stop using pull ups or nappies.
- Encourage regular use of the toilet 4-7 times a day.
- An extra wee before bedtime boys and girls should sit to wee.
- Use of a footstool or step.
- Keep on top of constipation, any pain, strain or discomfort visit your GP.
- Understand that constipation does affect wetting, it needs to be resolved before the child can become fully dry at night.

Plenty of praise for every small step

If your child does not wake up to the alarm:

- Wake child up fully turning lights on if necessary.
- Encourage child to turn off alarm.
- Take to the toilet and sit to wee.
- Your child should help you change the bed and night clothes.

Do not worry if this is the only progress you make to begin with.



Remember plenty of praise for every small step



The alarm is a training device

- 1. Child will begin to wake when the alarm sounds.
- 2. Eventually waking when their bladder is full.
- 3. Wake before the alarm goes off.
- 4. This process can take between 3 6 months.

It is important to have patience.

It is important to give praise.

It is very important not to blame the child.

Our aim is: to hold a telephone review after 4 weeks of using the alarm, to discuss in detail how your child is progressing and make a plan for the next two months.

This will then be followed with a further telephone appointment three months later, if not before.

We will provide you with charts to help record drinks and alarm progress.

Please feel free to contact the Continence Clinic at any time.

Further information and advice for both children and parents can be found at: <u>www.eric.org.uk</u>

Bedwetting alarms your questions answered.

A supplement to ERIC's Guide to Night Time Wetting.

er

